


---

# Linear Algebra notes

---

*Author*  
GABRIEL HEAROT

March 25, 2020





# Contents

<b>1 Basic Notions.</b>	<b>5</b>
1.1 Vector spaces . . . . .	5
1.2 Linear combinations . . . . .	5



# Chapter 1

## Basic Notions.

### 1.1 Vector spaces

**Definition** A vector space  $V$  is an object made of two sets: the set of the vectors, with its elements being denoted by a lowercase bold letter (e.g.  $\mathbf{u}$ ), and the set of the scalars, usually denoted by a Greek letter.  $\mathbf{u}$  and  $\mathbf{v}$  usually denote general vectors, while  $\alpha$  and  $\beta$  general scalars.

There are eight properties which vectors hold:

1. Commutativity:  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u} \quad \forall \mathbf{u}, \mathbf{v} \in V$ ;
2. Associativity:  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w}) \quad \forall \mathbf{u}, \mathbf{v}, \mathbf{w} \in V$ ;
3. Zero vector: there exists a unique vector  $\mathbf{0}$  such that  $\mathbf{v} + \mathbf{0} = \mathbf{v} \quad \forall \mathbf{v} \in V$ ;
4. Inverse vector: there always exists a vector  $-\mathbf{v}$  such that  $\mathbf{v} + (-\mathbf{v}) = \mathbf{v} - \mathbf{v} = \mathbf{0} \quad \forall \mathbf{v} \in V$ ;
5. Neutral scalar for multiplication: there always exists a scalar 1 such that  $1\mathbf{v} = \mathbf{v} \quad \forall \mathbf{v} \in V$ ;
6. Multiplicative associativity:  $(\alpha\beta)\mathbf{v} = \alpha(\beta\mathbf{v}) \quad \forall \mathbf{v} \in V$  and  $\forall \alpha, \beta$  which are scalars;
7. Distributive property:  $\alpha(\mathbf{u} + \mathbf{v}) = \alpha\mathbf{u} + \alpha\mathbf{v} \quad \forall \mathbf{u}, \mathbf{v} \in V$  and  $\forall \alpha$  which is scalar;
8. Vectorial distributive property:  $(\alpha + \beta)\mathbf{v} = \alpha\mathbf{v} + \beta\mathbf{v} \quad \forall \mathbf{v} \in V$  and  $\forall \alpha, \beta$  which are scalars.

If a vector space is made of scalars which are real numbers, it is called a *real* vector space. Likewise, for complex numbers, it is called a *complex* vector space.

**Matrices & polynomials** A space  $M_{m \times n}$  (sometimes written as  $M_{m,n}$ ) denotes a vectorial space made of  $m \times n$  matrices. A space  $\mathbb{P}_n$  denotes a vectorial space made of  $n$  degree polynomials.

A matrix  $A$  is an array of  $m$  rows and  $n$  columns and it is written as 
$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ \vdots & \ddots & \ddots & a_{2n} \\ a_{m1} & \dots & \dots & a_{mn} \end{pmatrix}.$$

$A^T$  represents the transposed matrix (i.e. a matrix  $A$  for which the rows & columns have been inverted).

### 1.2 Linear combinations

**Definition** Given a collection of vectors  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4, \dots, \mathbf{v}_n$  and a collection of scalars  $\alpha_1, \alpha_2, \alpha_3, \alpha_4, \dots, \alpha_n$ , the sum  $\sum_{p=1}^n \alpha_p \mathbf{v}_p$  is called a *linear combination*.

If there exists a unique representation for the *linear combination* of a set of vectors such that  $\forall \mathbf{v} \in V, \mathbf{v} = \sum_{p=1}^n \alpha_p \mathbf{v}_p$ , these ones are called a *basis*, while their coefficients are called *coordinates*.

A set of vectors is a *basis* if  $\mathbf{v} = \sum_{p=1}^n x_p \mathbf{v}_p$  admits a unique set of solutions.